

DOI: 10.2478/ffp-2019-0004

## Fungi isolated from shoots showing ash dieback in the Wolica Nature Reserve in Poland and artificially inoculated seedlings with *Hymenoscyphus fraxineus*

Anna Żółciak<sup>1</sup> ✉, Justyna Anna Nowakowska<sup>2</sup>, Artur Pacia<sup>3</sup>,  
Nenad Keča<sup>4</sup>, Tomasz Oszako<sup>5</sup>

<sup>1</sup> Forest Research Institute, Department of Forest Protection, Sękocin Stary, Braci Leśnej 3, 05-090 Raszyn, Poland, phone: +48 227153822, email: A.Zolciak@ibles.waw.pl

<sup>2</sup> Cardinal Stefan Wyszyński University in Warsaw, Faculty of Biology and Environmental Sciences, Wóycickiego 1/3, 01-938 Warsaw, Poland

<sup>3</sup> Forest District Chojnów, Piława, Klonowa 13, 05-532 Baniocha, Poland

<sup>4</sup> University of Belgrade, Faculty of Forestry, Kneza Višeslava 1, 11030 Belgrade, Serbia

<sup>5</sup> Białystok University of Technology, Faculty of Forestry in Hajnówka, Piłsudskiego 8, 17-200 Hajnówka, Poland

### ABSTRACT

Ash dieback caused by an alien, invasive fungus *Hymenoscyphus fraxineus* is a serious disease of European ash species in many parts in Europe. In Poland, the disease was recorded in the beginning of the 1990s. This study was performed in 2016–2017 with the aim to identify fungi isolated from ash shoots showing dieback symptoms in the Wolica Nature Reserve in Poland, as well as from shoots of two years-old ash seedlings inoculated with *H. fraxineus* in the greenhouse.

The most frequently isolated fungi from shoots of common ash (associated with the pathogenic fungus *H. fraxineus*) were identified on the basis of sequencing of the internal transcribed spacer region (ITS1) of fungal rDNA. In total, 19 fungal taxa were identified for ash shoots as follows: *H. fraxineus*, *Fusarium avenaceum*, *Alternaria* spp., *Phomopsis oblonga*, *Diplodia mutila* and other *Phomopsis* spp. The pathogen *H. fraxineus* was not found for all the shoots samples; one year after inoculation the aforementioned fungi and other species as: *Alternaria alternata*, *Bionectria ochroleuca*, *Epicoccum nigrum*, *F. acuminatum*, *F. avenaceum*, and *Paraphaesphaeria neglecta* were identified in inoculation point, as well as *H. fraxineus*. The same quantitative and qualitative changes of organisms were observed in the case of artificially colonised ash seedlings in the greenhouse, as well as in the shoots of adult ash trees in the forest.

### KEY WORDS

bark necrosis, ash dieback, *Fraxinus excelsior*, *Hymenoscyphus fraxineus*, ITS1-F

## INTRODUCTION

The dieback symptoms of ash have been observed in Poland since 1992 (Stocki and Stocka 1999; Sierota 2001; Przybył 2002; Gil et al. 2017). Since then, it is known that the fungus *Hymenoscyphus fraxineus* (T. Kowalski) Baral, Queloz and Hosoya is responsible for a severe decline on all age classes of *Fraxinus excelsior* L. stands (Kowalski and Holdenrieder 2008, 2009; Kirisits et al. 2010). Ash dieback caused by *H. fraxineus* has been reported in numerous countries in Europe (Vasaitis and Enderle 2017; Enderle et al. 2017).

It is well documented that after the fungus *H. fraxineus* colonization of bark and wood tissues, phytotoxin viridiol is secreted causing dying of plant tissues (Andersson et al. 2010). After primary intensive development and killing of bark tissues, pathogen moves to the wood where it could be quickly substituted with other colonizers of dead tissues (Kowalski et al. 2016). Several species of fungi were found in necrotic tissues of shoots and stems of dying common ash trees (Przybył 2002; Kowalski and Kehr 1992; Kowalski and Łukomska 2005; Schumacher et al. 2007; Bakys et al. 2009; Kowalski and Bartnik 2010). Species observed on dead tissue of ash shoots represent a cosmopolitan fungal genus *Alternaria* (that includes saprobic and pathogenic species associated with a wide variety of substrates) (Woudenberg et al. 2013; Kowalski et al. 2016), genus *Epicoccum* (saprophytes or a weak pathogens or endophytes) (Khulbe et al. 1992; Bruton et al. 1993; Fávoro et al. 2012; Ristić et al. 2012), *Diaporthe/Phomopsis* (that includes pathogens, endophytes and saprobes) (Udayanga et al. 2011; Gomes et al. 2013; Kowalski et al. 2016), a genus *Fusarium* (that includes pathogens, but most species of this genus are harmless saprobes) (Summerell et al. 2010; Kowalski et al. 2016; Kowalski et al. 2017).

Study of the diversity of fungi in natural reserves is of wider interest because it is well known that more diverse ecosystems better cope with disturbing factors and alien invasive species. Very often, the molecular methods are applied in order to determine the fungal species with the help of universal primers, that is, 'ITS1' and 'ITS4' (Gardes and Bruns 1993; <https://www.ncbi.nlm.nih.gov/>). The above primers are complementary to the Internal Transcribed Sequences (ITS), and gained

wide acceptance for work with fungal identification, because they amplify the highly variable ITS1 and ITS2 regions surrounding the 5.8S-coding sequence situated between the Small SubUnit-coding sequence (SSU) and the Large SubUnit-coding sequence (LSU) of the ribosomal operon (Martin and Rygiewicz 2005; Porras-Alfaro et al. 2014). Because of not exclusive power of distinction between fungal and plant rDNA amplification provided by 'ITS1' primer, the more accurate primer for fungal identification ('ITS1-F', Gardes and Bruns 1993), besides the 'ITS1' primer has been used in the present study.

The aim of this study was to identify fungal species associated with necrotic tissues of shoots of dying ash in the Wolica Nature Reserve in Poland and to compare results with data obtained after artificial inoculations of two-year-old ash seedlings with *H. fraxineus*.

## MATERIAL AND METHODS

### Experiment I

#### Sampling in the stand

In spring 2016, about thirty shoots, 4–8 mm in diameter with necrotic lesions were sampled from three common ash trees (*F. excelsior*) in the 26-year-old ash stand. The stand is located in compartment 374 c in the Wolica Nature Reserve (52°11'16.70"N, 20°41'17.96"E). The phenomenon of ash dieback was investigated in this stand in 2012 and 2013, when the vitality of trees was very poor and many trees showed severe defoliation (Pacia et al. 2017).

In order to produce inocula for artificial inoculation, some fresh and healthy looking ash shoots were collected from mixed stands in the Chojnów Forest District.

#### Isolation and identification of microorganisms from ash trees

In the spring season in 2016, the living, symptomatic shoots (with lesions) collected in Wolica ash stand were cut into pieces of the length ca. 1–2 cm, and then sterilized in 90% ethanol (1 min), 4% NaOCl sodium hypochlorite (5 min) and again in 90% ethanol (30 s) (Kowalski 2006). After drying them on a sterile filter paper and cutting down the top layer of bark, the wood was cut for pieces of ca. 5 × 2 × 2 mm and put on 2%

malt-extract agar (MEA) medium (Merck, Germany) solidified in Petri plates. It was then incubated at 20°C in dark. In total, 60 pieces of symptomatic wood tissues were taken from thirty shoots.

The Petri plates with inoculum were incubated at 20°C in the dark and few days old growing colonies were transferred onto a new MEA media. The pure cultures of isolated fungi (22) were stored in Forest Research Institute and used for DNA extraction.

The 50 mg of mycelium of each fungal species were homogenized in the presence of liquid nitrogen using a mortar and pestle. Total genomic DNA was extracted from mycelium with DNeasy Plant mini kit (Qiagen®, Hilden, Germany) according to the manufacturer's instructions.

In order to identify the fungal species, the DNA of 22 fungal isolates were amplified by a conventional PCR approach with the use of the universal primers ITS1 (5' -TCC GTA GGT GAA CCT GCG G, White et al. 1990) or ITS1-F (5' -CTT GGT CAT TTA GAG GAA GTA A, Gardes and Bruns 1993) were used (Tab. 1). The 25 µl PCR mix consisted of PCR buffer (10x), 25 mM MgCl<sub>2</sub>, 0.2 mM dNTPs 1 µM each primer, 1 U of Taq polymerase, MQ water up to 25 µl and 100 ng of DNA. Cycling conditions were denaturation 95°C 5 min followed by 35 cycles of denaturation 94°C for 0.5 min, annealing at 56°C for 0.5 min and extension 72°C 1 min, and a final extension at 72°C for 7 min (Korabecna 2007, modified by Nowakowska). The PCR products of ITS region were separated by electrophore-

**Table 1.** Identification of 22 fungal isolates from dead tissues of ash shoots by using rDNA ITS sequencing

Species isolated	Primers used for PCR amplification	Reference number in NCBI database and similarity (%)	No. of accession in Genbank
<i>Alternaria alternata</i> (Fr.) Keissl. <sup>1</sup>	ITS1-F	KY609180 (99.0)	MF509751
<i>Alternaria</i> sp. <sup>2</sup>	ITS1-F	EU366278 (99.0)	MF509752
<i>Bionectria ochroleuca</i> (Schwein.) Schroers & Samuels <sup>2</sup>	ITS1-F	GU566253 (100.0)	MF509749
<i>Boeremia exigua</i> (Desm.) Aveskamp, Gruyter & Verkley <sup>1</sup>	ITS1	MF599108 (100.0)	KX618484
<i>Boeremia exigua</i> var. <i>exigua</i> <sup>1</sup>	ITS1	MF435055 (99.0)	KX618485
			KX618486
<i>Cryptosphaeria eunomia</i> (Fr.) Fuckel <sup>1</sup>	ITS1	KY613993 (100.0)	KX618490
<i>Cryptosphaeria eunomia</i> <sup>1</sup>	ITS1	KY613993 (100.0)	KX618491
<i>Diaporthe eres</i> Nitschke <sup>2</sup>	ITS1-F	EU571099 (198.0)	MF509750
<i>Diplodia mutila</i> (Fr.) Mont. <sup>1</sup>	ITS1	KF766158 (100.0)	KX618487
<i>Diplodia mutila</i> (Fr.) Mont. <sup>1</sup>	ITS1	KF766158 (100.0)	KX618488
<i>Epicoccum nigrum</i> Link <sup>2</sup>	ITS1-F	KR094452 (100.0)	MF509753
<i>Fusarium acuminatum</i> Ellis & Everh. <sup>2</sup>	ITS1-F	KJ737377 (100.0)	MF509746
<i>Fusarium avenaceum</i> (Fr.) Sacc. <sup>2</sup>	ITS1-F	HG936695 (99.0)	MF509747
<i>Fusarium oxysporum</i> Schlechtend. <sup>1</sup>	ITS1	AB470850 (100.0)	KX618492
<i>Hymenoscyphus fraxineus</i> Baral. et al. <sup>1,*</sup>	ITS1	HM193468.1 (100.0)	KY613994
<i>Paraphaesphaeria neglecta</i> Verkley, Riccioni & Stielow <sup>2</sup>	ITS1-F	JX496164 (100.0)	MF509754
<i>Phomopsis oblonga</i> (Desm.) Traverso <sup>1</sup>	ITS1	KR909214.1 (100.0)	KX618489
<i>Phomopsis</i> sp. <sup>2</sup>	ITS1-F	EU571102 (99.0)	MF509748
Unidentified <i>Ascomycota</i> <sup>1</sup>	ITS1	JQ765657 (99.0)	KX618493
Unidentified <i>Ascomycota</i> <sup>2</sup>	ITS1	JQ765657 (99.0)	KX618494
Unidentified <i>Pleosporales</i> <sup>1</sup>	ITS1	KC311498 (99.0)	KX618495

1 – isolated from dying ash shoots taken from the ash trees in the Wolica Nature Reserve; 2 – isolated from 2-year-old dead ash seedlings one year after the inoculation of *H. fraxineus*.

\* Under the rules for the naming of fungi with pleomorphic life-cycles adopted in 2011, the nomenclaturally correct name for the fungus causing ash dieback in Europe is determined to be *Hymenoscyphus fraxineus*, with the basionym *Chalara fraxinea*.

sis on 1.5% agarose gels. Prior to sequencing, the amplicons were purified with QIAquick PCR Purification Kit (Qiagen Ltd), according to the manufacturer's protocol. Sequencing was performed using the BigDye Terminal Cycle Sequencing Kit (AB Applied Biosystems, CA, USA) in ABI 3500 Genetic Analyzer (Life Technologies™, USA). The retrieved sequences were identified by comparison with sequences deposited in NCBI database using BLAST (Basic Local Alignment Search Tool, accessed November 8, 2017).

All the selected sequences based on the analyses of the ITS1 rDNA region were compared by aligning them with an online version of Multiple Sequence Alignment (MSA) with Clustal Omega program v. 1.2.4 (Sievers et al. 2011). To better illustrate the relationship between the identified isolates, the phylogenetic Neighbour-Joining tree has been constructed in MAFFT v. 7 with bootstrap 100, threshold  $8.4e^{-11}$  (<https://mafft.cbrc.jp/alignment/server/>).

## Experiment II

### Plant material

The experiment was performed in the greenhouse of the Forest Research Institute (IBL, Sękocin Stary, Poland). Sixty two-year-old European ash seedlings were planted in 1 l pots filled with a 1:1 (v:v) peat: perlite mixture at the beginning of the vegetation in May 2016. The temperature ca. 22°C and relative humidity  $65\% \pm 5\%$  was controlled automatically by the computer software and were shaded when the light intensity exceeded 50 klx. Plants were irrigated manually with tap water once a day.

### Inoculation of seedlings

In September 2016, twenty ash seedlings were inoculated at the base of stems (5–10 cm above the soil level). The control variant consisted of forty ash seedlings: twenty wounded in the same way as described above and another twenty ones not inoculated and not wounded. In order to obtain the inocula, the healthy ash branches were cut into small pieces autoclaved and placed on the surface of the *H. fraxineus* four-week-old pure cultures and kept for 4 weeks to let them overgrow with fungus. Then, the inocula were placed under the cut bark in T shape and sealed with Parafilm (Sigma-Aldrich, Taufkirchen, Germany).


### Isolation and identification of microorganisms from seedlings

In September 2017 (one year after inoculation), the re-isolation from 20 dead ash seedlings inoculated with *H. fraxineus* were performed from around the inoculation points and 120 pieces of wood were placed on 2% malt-extract agar (MEA) medium (Merck, Germany). The obtained pure cultures of fungi were identified in both ways morphologically at the genus level and molecularly (according to the method described in the Experiment I) at the level of species.

## RESULTS

From shoots of ash trees in Wolica stand, the following fungi species were identified: *Phomopsis oblonga* (21.7% of all), *Diplodia mutila* (20.5%), *Boeremia exigua* syn. *Boeremia exigua* var. *exigua* (20.9%), *Cryptosphaeria eunomia* (8.6%), *Fusarium oxysporum* (1.6%) and *Ascomycota* sp. (13.8%), *Pleosporales* sp. (8.3%) and *Hymenoscyphus fraxinues* (4.6%) (Tab. 1). The phylogenetic alignment based on the NJ tree confirmed the close relationships between ITS1 rDNA sequences of *Fusarium* isolates, and, for example, *Fusarium* sp. (accession numbers MF509746, MF50974647 and KX618492), and *Bionectria ochroleuca* (MF5097749) (Fig. 1). The latest four isolates were characterised by very low genetic distance (0.038). Polymorphic sites revealed between positions 0–994 bp of the studied ITS1 regions clearly shows the differences between all the investigated species (Suppl. fig. in electronic ver.).

In the experiment in greenhouse, 100% of the seedlings inoculated with *H. fraxineus* were dead (already eight months after inoculation). From ash seedlings, *H. fraxineus* was identified in only 2.4% of the obtained cultures. Among the obtained isolates dominated, *Fusarium* species were dominated with 38.8% (*F. avenaceum* – 33.1% and *F. acuminatum* – 5.5%), while *Alternaria* species with the frequency of 28.7%. *Phomopsis* spp., including *Diplodia eres*, were third most frequently observed colonizers with the frequency of 17.7%, while the other unidentified species were from phylum *Ascomycota* participated with 7.9% in total isolations. *Bionectria ochroleuca*, *Paraphaesaeria neglecta* and *E. nigrum* were observed at frequencies 4%, 0.4 and 0.1%, respectively.



**Figure 1.** NJ tree derived from the analysis of the ITS dataset. For each isolate, the number of accession to the GenBank and the distance representing the nucleotide substitution values were given

## DISCUSSION

Do the present studies show the succession of fungi colonizing ash tissues? We did not apply different time of sampling (initial necrosis, advanced necrosis and dead shoots); however, we noticed a clear difference after a year of the experiment. Instead of inoculated pathogenic anamorph *Chalara fraxinea*, we isolated many other fungal species. The reason for this is probably that the fungus *H. fraxineus* is a primary pathogen killing healthy tissues, mainly to its ability of viridol production, which is toxic to ash plants (Grad et al. 2009; Andersson et al. 2010). However, later, the fungus is losing the competition with other decomposing fungi being more successful and taking over their domination in dead tissues. We noticed that symptomatic ash shoots colonized by other plant pathogens, as well as by endophytes and saprobes.

In this work, the fungal species were identified based on the analysis of ITS1 sequences of the rDNA. Identification of fungi from, for example, *Fusarium* and *Alternaria* genera based only on analysis of ITS sequences are very problematic, nevertheless the ITS1 region is commonly used as the first step in molecular identification of fungi (Toju et al. 2012). It has been demonstrated that this fragment of internal transcribed spacer of fungal ribosomal RNA identifies the ascomycetous, basidiomycetous, and zygomycetous fungi to the species level (Glass and Donaldson 1995). Other genes (e.g., ACT,  $\beta$ T, TEF1- $\alpha$ , CAL) can also be studied for a precise identification of fungal species. In many studies, the use of ITS sequence data together with protein-coding genetic data has become the normal practice for fungal identification (Korabecna 2007; Duong et al. 2012). According to Porrás-Alfaro et al. (2014), the sequencing of the LSU region in many fungal spe-

cies gives very similar results of the comparison based on MSA with the study based on the entire ITS region (ITS1 plus 5.8S rRNA plus ITS2). In some cases, the phylogenetic analyses of the ITS2-LSU sequences have revealed that the morphologically similar isolates represent a complex of the cryptic species (Doung et al. 2012). Hence, our molecular analysis with the general set of primers, that is, ITS1 and ITS1-F can be considered as a preliminary attempt of identification, prior to the future focus on other genes, that is, ACT,  $\beta$ -T, CAL or TEF1- $\alpha$ , which permit more accurate identification of fungi from, for example, *Fusarium* and *Alternaria* genera.

Nevertheless, we managed to describe the fungal community structure in the declining shoots of ash trees in the Wolica Nature Reserve in Poland and we got similar results in the greenhouse experiment, while inoculating shoots with *H. fraxineus* and performing re-isolation tests after a year. Based on the comparison of ITS1 sequences, the results of this survey revealed 19 fungal taxa associated with ash shoots showing dieback. This is the valuable point of this research. Moreover, our findings confirmed the occurrence of fungi in symptomatic shoots of *F. excelsior* published earlier by many scientists. According to Boudier (1994), *Phoma exigua* (*Boeremia exigua*) can cause brown spotting on 1- and 2-year-old shoots of European ash seedling. Next to *Hysteroglyphium fraxini*, *P. exigua* have been considered to be involved in serious decline, in combination with abiotic factors (Grzywacz 1995; Przybył 2001). *Boeremia exigua* (synonym *B. exigua* var. *exigua*) is an ubiquitous soil borne saprobe, weak pathogen or wound parasite. Associated with stem and leaf lesions of a wide range of host plants and with rotting fleshy roots and tubers, often causing distinct symptoms such as leaf spots, stem lesions, damping off, dieback, root rots and tuber rots. Our findings are also compatible with Kowalski and Kehr (1992) research, who presented population of endophytic fungi on ash branches with the most characteristic fungi from *Coniothyrium*, *Pezicula* and *Chalara* genus. They found the same species of fungi as we did like *Alternaria alternata*, *Diplodia mutila*, and *Phomopsis* sp., but also *Cladosporium cladosporioides*, *Cytospora ambiens* and *Phomopsis controversa*. The species *Phomopsis scobina* was isolated from dead or dying top parts of shoots (Kowalski and Łukomska 2005), while from cankers also *C. am-*

*biens*, *P. controversa*, and *P. scobina*. Przybył (2001) noticed that *P. controversa*, *P. scobina*, *C. ambiens*, *A. alternata* and *D. mutila* are the fungi associated with both dead and living 1–2-year-old trees showing necrosis of their tops, which was also true in our case for *A. alternata* and *D. mutila*.

Fungi belonging to *Phomopsis* and *Cytospora* genus are pathogens of weakness and are often present in dying shoots (Przybył 2002; Cech 2005; Kowalski and Łukomska 2005; Barklund 2006; Gomes et al. 2013). *D. mutila* (telemorph: *Botryosphaeria stevensii*) is a widespread pathogen in the world occurring mainly on *Quercus*, *Juniperus*, and *Pinus* (Vajna 1986; Luque and Girbal 1989; El-Badri et al. 1998; Stanosz et al. 1998; Mohali and Encinas 2001). Aforementioned fungi can exist in oak shoots as endophytes, but can also become pathogenic when the host is weakened by abiotic factors (Collado et al. 1999; Ragazzi et al. 2001). Przybył (2003) and Kowalski et al. (2017) proved pathogenicity of *D. mutila*. According to Kowalski et al. (2017), *D. mutila* and *Cytospora pruinosa* are less pathogenic than *H. fraxineus*, but more than the other fungi. Species like *Diaporthe eres*, *F. avenaceum*, *F. solani* and *F. lateritium* are considered to be the least pathogenic.

The occurrence of *Epicoccum* is also not so simple and obvious case in respect of its pathogenicity. Namely, this genus could be a saprophyte, a weak pathogen of rapeseed seeds and roots as well as sorghum seeds and seedlings, an opportunistic pathogen of melon, cucumber, tomato, apple and pear fruit or sugarcane endophyte (Khulbe et al. 1992). Seed-borne infection, pathogenic importance and control of *Epicoccum nigrum* Ehrenb, ex Schlecht. In rape seed (*Brassica napus* L.), these were thoroughly described by Khulbe et al. (1992), Bruton et al. (1993), Fávoro et al. (2012) and Ristić et al. (2012).

Tulik et al. (2017) developed the hypothesis that ash trees affected by *H. fraxineus* causing ash dieback could exhibit a reduced vessel size and density along the trunk, as well as a lowered width of annual wood rings, leading to the weakening of water transport towards the crown. As the consequence, shoots that are dying result in infection by aforementioned species of fungi genus *Phomopsis*, *Diplodia*, *Fusicoccum*, *Diaporthe*, *Cytospora* and others. In such a way, the phenomenon of fungi succession is initiated first in living, later in dead ash shoots. This observation proved to be a rapid

stepping down of *H. fraxineus* – the primary cause of ash dieback, being replaced by other fungi colonizing ash tissues.

From this point of view, the knowledge of the distribution of pathogenic fungal species associated with *F. excelsior* suffering from ash-dieback phenomenon can be very useful in forest tree management and diversity conservation programs. In both experiments (in the stand and in the greenhouse), we observed no crucial difference between the fungal species isolated from young shoots (naturally infected in the ash stand and seedlings inoculated with *H. fraxineus* in the greenhouse). However, we did not check the occurrence of the aforementioned fungi in living plant tissues. We believed that those fungi come after infection by *H. fraxineus* because control plants were healthy. Nevertheless, it is possible that some of those fungi could live as endophytes, but it was not the purpose of this study.

## CONCLUSION

After ash shoot infection by *H. fraxineus*, the succession of other species of fungi was observed as well in natural (Wolica Reserve) as in control conditions (greenhouse).

Especially in fungi, which are less successful in colonizing living tissues, start to overtake wood of ash shoots previously killed by *H. fraxineus*.

## ACKNOWLEDGEMENTS

Authors thank Małgorzata Gorzkowska for the ITS1 rDNA analysis, and IBL funds supporting 3 months stay of Prof. Nenad Keča in the Department of Forest Protection in Sękocin Stary.

The present research was supported by the Ministry of Science and Higher Education of Poland as the part of project no. 240 325.

## REFERENCES

Andersson, P.F., Johansson, S.B.K., Stenlid, J., Broberg, A. 2010. Isolation, identification and necrotic activity of viridiol from *Chalara fraxinea*, the fungus

- responsible for dieback of ash. *Forest Pathology*, 40, 43–46. DOI: 10.1111/j.1439-0329.2009.00605.x
- Bakys, R., Vasaitis, R., Barklund, P., Thomsen, I.M., Stenlid, J. 2009. Occurrence and pathogenicity of fungi in necrotic and non-symptomatic shoots of declining common ash (*Fraxinus excelsior*) in Sweden. *European Journal of Forest Research*, 128, 51–60.
- Barklund, P. 2006. Unknown fungus causes dieback of ash shoots (in Swedish). *Skogseko*, 3, 10–11.
- Boudier, B. 1994. The canker of ash seedlings (in French). *Phytoma, La Défense des Végétaux*, 461, 35–36.
- Bruton, B.D., Redlin, S.C., Collins, J.K., Sams, C.E. 1993. Postharvest decay of cantaloupe caused by *Epicoccum nigrum*. *Plant Disease*, 77, 1060–1062.
- Cech, L.T. 2006. Auffelende Schadfaktoren an Waldbaeumen im Jahr 2005. *Forstschutz Aktuell*, 35, 6–7.
- Collado, J., Platas, G., Gonzalews, I., Pelaez, F. 1999. Geographical and seasonal influences on the distribution of fungal endophytes in *Quercus ilex*. *New Phytologist*, 144, 525–532.
- Duong, T.A., De Beer, Z.W., Wingfield, B.D., Wingfield, M.J. 2012. Phylogeny and taxonomy of species in the *Grosmannia serpens* complex. *Mycologia*, 104, 715–732.
- El-Badri, N., Abadie, M. 1999. Contribution to the ultrastructural study of *Diplodia mutila* (Fri.) Mont. implicated in the decline process of the cork oak *Quercus suber* L., in Morocco. *IOBC/WPRS Bulletin*, 22 (3), 13–18.
- Enderle, R., Pliura, A., Vasaitis, R. (eds.). 2017. Advances in ash dieback research, – and some other invasive diseases of trees. *Baltic Forestry*, 23 (1), 1–333.
- Fávaro, L.C. de Lima, Sebastianes, F.L. de Souza, Araújo, W.L. 2012. *Epicoccum nigrum* P16, a sugarcane endophyte, produces antifungal compounds and induces root growth. *PLoS ONE*, 7, e36826. DOI: 10.1371/journal.pone.0036826
- Gardes, M., Bruns, T.D. 1993. ITS primers with enhanced specificity for basidiomycetes application to the identification of mycorrhizae and rusts. *Molecular Ecology*, 2, 113–118. DOI: 10.1111/j.1365-294X.1993.tb00005.x
- Gil, W., Kowalski, T., Kraj, W., Zachara, T., Łukasiewicz, J., Paluch, R., Nowakowska, J.A., Oszako, T.

2017. Ash dieback in Poland – history of the phenomenon and possibilities of its limitation. In: Dieback of European ash (*Fraxinus* spp.). Consequences and guidelines for sustainable management (eds.: R. Vasaitis, R. Enderle), Swedish University of Agricultural Sciences, Uppsala, Sweden, 176–184.
- Glass, N.L., Donaldson, G.C. 1995. Development of primer sets designed for use with the PCR to amplify conserved genes from filamentous ascomycetes. *Applied Environmental Microbiology*, 61, 1323–1330.
- Gomes, R.R., Glienke, C., Videira, S.I.R., Lombard, L., Groenewald, J.Z., Crous, P.W. 2013. *Diaporthe*: a genus of endophytic, saprobic and plant pathogenic fungi. *Persoonia*, 31, 1–41. DOI: 10.3767/003158513X666844
- Grad, B., Kowalski, T., Kraj, W. 2009. Studies on secondary metabolite produced by *Chalara fraxinea* and its phytotoxic influence on *Fraxinus excelsior*. *Phytopathologia Polonica*, 54, 61–69.
- Grzywacz, A. 1995. Ważniejsze choroby infekcyjne. In: Jesion wyniosły *Fraxinus excelsior* L. (ed.: W. Bugała.). Wyd. Sorus, Poznań-Kórnik, 371–415.
- Khulbe, R.D., Dhyani, A.P., Sati, M.C. 1992. Seed-borne Infection, pathogenic importance and control of *Epicoccum nigrum* Ehrenb, ex Schlecht. in rape seed (*Brassica napus* L.) in Kumaun Hills, U.P., India. *Proceedings of the Indian National Science Academy*, B58 (2/3), 135–140.
- Kirisits, T., Matlakova, M., Mottinger-Kroupa, S., Cech, T.L., Halmschlager, E. 2010. *Chalara fraxinea* associated with dieback of narrow-leaved ash (*Fraxinus angustifolia*). *Plant Pathology*, 59, 411.
- Korabecna, M. 2007. The variability in the fungal ribosomal DNA (ITS1, ITS2, and 5.8 S rRNA Gene): its biological meaning and application in medical mycology. *Communicating Current Research and Educational Topics and Trends in Applied Microbiology*, 105, 783–787.
- Kowalski, T. 2006. *Chalara fraxinea* sp. nov. associated with dieback of ash (*Fraxinus excelsior*) in Poland. *Forest Pathology*, 36, 264–270. DOI: 10.1111/j.1439-0329.2006.00453.x
- Kowalski, T., Bartnik, C. 2010. Morphological variation in colonies of *Chalara fraxinea* isolated from ash (*Fraxinus excelsior* L.) stems with symptoms of dieback and effects of temperature on colony growth and structure. *Acta Agrobotanica*, 63, 99–106.
- Kowalski, T., Bilański, P., Kraj, W. 2017. Pathogenicity of fungi associated with ash dieback towards *Fraxinus excelsior*. *Plant Pathology*, 66 (8), 1228–1238. DOI: 10.1111/ppa.12667
- Kowalski, T., Holdenrieder, O. 2008. Eine neue Pilzkrankheit an Esche in Europa. *Schweizerische Zeitschrift für Forstwesen*, 3, 45–50.
- Kowalski, T., Holdenrieder, O. 2009. Pathogenicity of *Chalara fraxinea*. *Forest Pathology*, 39, 1–7.
- Kowalski, T., Kehr, R.D. 1992. Endophytic fungal colonization of branch bases in several forest tree species. *Sydowia*, 44, 137–168.
- Kowalski, T., Kraj, W., Bednarz, B. 2016. Fungi on stems and twigs in initial and advanced stages of dieback of European ash (*Fraxinus excelsior*) in Poland. *European Journal of Forest Research*, 135, 565–579.
- Kowalski, T., Łukomska, A. 2005. Badania nad zamierniem jesionu (*Fraxinus excelsior* L.) w drzewostanach Nadleśnictwa Włoszczowa. *Acta Agrobotanica*, 59, 429–440.
- Luque, J., Girbal, J. 1989. Dieback of cork oak (*Quercus suber*) in Catalonia (NE Spain) caused by *Botryosphaeria stevensii*. *European Journal of Forest Pathology*, 19, 7–13.
- Martin, K.J., Rygiewicz, P.T. 2005. Fungal-specific PCR primers developed for analysis of the ITS region of environmental DNA extracts. *BMC Microbiology*, 5 (1), 28. DOI: 10.1186/1471-2180-5-28
- Mohali, S., Encinas, O. 2001. Association of *Diplodia mutila* with blue stain of Caribbean pine in Venezuela. *Forest Pathology*, 31, 187–189.
- Pacia, A., Nowakowska, J.A., Tkaczyk, M., Sikora, K., Tereba, A., Borys, M., Milenković, I., Pszczółkowska, A., Okorski, A., Oszako, T. 2017. Common ash stand affected by ash dieback in the Wolica Nature Reserve in Poland. *Baltic Forestry*, 23(1), 183–197.
- Porras-Alfaro, A., Liu, K.L., Kuske, C.R., Xie, G. 2014. From genus to phylum: large-subunit and internal transcribed spacer rRNA operon regions show similar classification accuracies influenced by database composition. *Applied and Environmental Microbiology*, 80 (3), 829–840.
- Przybył, K. 2001. Grzyby występujące w wierzchołkowej części pędów jesionu wyniosłego wykazujących zmiany nekrotyczne. Materiały z V Konferencji



- Sekcji Chorób Roślin Drzewiastych Polskiego Towarzystwa Fitopatologicznego. Etiologia i objawy chorób grzybowych oraz ich występowanie i szkodliwość w ekosystemach leśnych. Poznań–Błażewko, 29 maja–1 czerwca 2001, 32–41.
- Przybył, K. 2002. Fungi associated with necrotic apical parts of *Fraxinus excelsior* shoots. *Forest Pathology*, 32, 387–394.
- Przybył, K. 2003. Effect of *Pseudomonas* spp. on inoculation of young plants of *Fraxinus excelsior* stem with *Diplodia mutila*. *Dendrobiology*, 50, 29–32.
- Ragazzi, A., Morrica, S., Capretti, P., Dellavalle, I., Mancini, F., Turco, E. 2001. Endophytic fungi in *Quercus cerris*: isolation frequency in relation to phenological phase, tree health and the organ affected. *Phytopathologia Mediterranea*, 40, 165–171.
- Ristić, D., Stanković, I., Vučurović, A., Berenji, J., Krnjajić, S., Krstić, B., Bulajić, A. 2012. *Epicoccum nigrum* novi patogen semena sirka u Srbiji. *Ratarstvo i povrtarstvo*, 49, 160–166.
- Schumacher, J., Wulf, A., Leonhard, S. 2007. Erster Nachweis von *Chalara fraxinea* T. Kowalski sp. nov. in Deutschland – ein Verursacher neuartiger Schäden an Eschen. *Nachrichtenblatt des Deutschen Pflanzenschutzdienstes*, 59, 121–123.
- Sierota, Z. 2001. Choroby lasu. Choroby lasu? Centrum Informacyjne Lasów Państwowych, 1–156. ISBN 83-88478-18-4.
- Sievers, F., Wilm, A., Dineen, D., Gibson, T.J., Karpus, K., Li, W., Lopez, R., Mc William, H., Remmert, M., Söding, J., Thompson, J.D., Higgins, D.G. 2011. Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. *Molecular Systems Biology*, 7, 539. DOI: 10.1038/msb.2011.75
- Stanosz, G.R., Swart, W.J., Smith, D.R. 1998. Similarity between fungi identified as *Diplodia pinea* f. sp. *cupressi* in Israel and *Botryosphaeria stevensii* or *Diplodia mutila* on *Juniperus* in the United States. *European Journal of Forest Pathology*, 28, 33–42.
- Stocki, J., Stocka, T. 1999. Przyczyny zamierania drzew i drzewostanów jesionowych. *Biblioteczka leśniczego*, 118. Wyd. Świat, Warszawa, 3–11.
- Summerell, B.A., Laurence, M.H., Liew, E.C., Leslie, J.F. 2010. Biogeography and phylogeography of *Fusarium*: a review. *Fungal Diversity*, 44, 3–13. DOI: 10.1007/s13225-010-0060-2
- Toju, H., Tanabe, A.S., Yamamoto, S., Sato, H. 2012. High-coverage ITS primers for the DNA-based identification of ascomycetes and basidiomycetes in environmental samples. *PLoS ONE*, 7, e40863.
- Tulik, M., Zakrzewski, J., Adamczyk, J., Tereba, A., Yaman, B., Nowakowska, J.A. 2017. Anatomical and genetic aspects of ash dieback: a look at the wood structure. *iForest-Biogeosciences and Forestry*, 10, 522–528. DOI: 10.3832/ifor2080-010
- Udayanga, D. 2011. The genus *Phomopsis*: biology, applications, species concepts and names of common phytopathogens. *Fungal Diversity*, 50, 189–225. DOI: 10.1007/s13225-011-0126-9
- Vajna, L. 1986. Branch canker and dieback of sessile oak (*Quercus petraea*) in Hungary caused by *Diplodia mutila*. *European Journal of Forest Pathology*, 16, 223–229.
- Vasaitis, R., Enderle, R. 2017. Dieback of European ash (*Fraxinus* spp.) – consequences and guidelines for sustainable management. The Report on European Cooperation in Science & Technology (COST). Action FP1103 FRAXBACK. Uppsala, 1–229.
- White, T.J., Bruns, T., Lee, S., Taylor, J. 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: PCR protocols: A guide to methods and applications (eds.: M.A. Innis et al.). Academic Press, New York, 315–322.
- Woudenberg, J.H., Groenewald, J.Z., Binder, M., Crous, P.W. 2013. *Alternaria* redefined. *Studies in Mycology*, 75, 171–212. DOI: 10.3114/sim0015

**Supplementary Figure.** Polymorphism sites in the ITS regions of the 21 investigated isolates. Numbers at the end of column indicate the relative positions in the alignments. Sites indicated by \* are conservative in all isolates, while sites which differ are indicated with ‘-’.

**CLUSTAL Omega(1.2.4) multiple sequence alignment**

Cryptosphaeria_eunomia_KX618490	-----	0
Cryptosphaeria_eunomia_KX618491	-----	0
Phomopsis_sp._MF509748	-----	0
Diaporthe_eres_MF509750	-----	0
Phomopsis_oblonga_KX618489	-----	0
Bionectria_ochroleuca_MF509749	-----	0
Fusarium_oxysporum_KX618492	-----	0
Fusarium_acumnatum_MF509746	-----	0
Fusarium_avenaceum_MF509747	-----	0
Hymenoscyphus_fraxineus_KY613994	-----	0
Diplodia_mutila_KX618487	-----	0
Diplodia_mutila_KX618488	-----	0
Paraphaeosphaeria_neglecta_MF509754	tcgccaagacacttgcgtaagccttgcagctggagacagtagcccaacgactgtaaaca	60
Alternaria_alternata_MF509751	-----	0
Alternaria_sp._MF509752	-----	0
Pleosporales_sp._KX618495	-----	0
Ascomycota_sp._KX618493	-----	0
Ascomycota_sp._KX618494	-----	0
Epicoccum_nigrum_MF509753	-----	0
Boeremia_exigua_KX618486	-----	0
Boeremia_exigua_KX618484	-----	0
Boeremia_exigua_KX618485	-----	0
Cryptosphaeria_eunomia_KX618490	-----	0
Cryptosphaeria_eunomia_KX618491	-----	0
Phomopsis_sp._MF509748	-----	0
Diaporthe_eres_MF509750	-----	0
Phomopsis_oblonga_KX618489	-----	0
Bionectria_ochroleuca_MF509749	-----	0
Fusarium_oxysporum_KX618492	-----	0
Fusarium_acumnatum_MF509746	-----	0
Fusarium_avenaceum_MF509747	-----	0
Hymenoscyphus_fraxineus_KY613994	-----	0
Diplodia_mutila_KX618487	-----	0
Diplodia_mutila_KX618488	-----	0
Paraphaeosphaeria_neglecta_MF509754	atggcgctcatgcaagtcagctccggctggcgacactttcgaattgcggggaaccctc	120
Alternaria_alternata_MF509751	-----	0
Alternaria_sp._MF509752	-----	0
Pleosporales_sp._KX618495	-----	0
Ascomycota_sp._KX618493	-----	0
Ascomycota_sp._KX618494	-----	0
Epicoccum_nigrum_MF509753	-----	0
Boeremia_exigua_KX618486	-----	0
Boeremia_exigua_KX618484	-----	0
Boeremia_exigua_KX618485	-----	0
Cryptosphaeria_eunomia_KX618490	-----	0
Cryptosphaeria_eunomia_KX618491	-----	0
Phomopsis_sp._MF509748	-----	0
Diaporthe_eres_MF509750	-----	0
Phomopsis_oblonga_KX618489	-----	0
Bionectria_ochroleuca_MF509749	-----	0
Fusarium_oxysporum_KX618492	-----	0
Fusarium_acumnatum_MF509746	-----	0
Fusarium_avenaceum_MF509747	-----	0
Hymenoscyphus_fraxineus_KY613994	-----	0
Diplodia_mutila_KX618487	-----	0
Diplodia_mutila_KX618488	-----	0
Paraphaeosphaeria_neglecta_MF509754	aagagctcttcacaccaagccagccgggaaaccggctggtggccattggaaaccatgg	180
Alternaria_alternata_MF509751	-----	0
Alternaria_sp._MF509752	-----	0
Pleosporales_sp._KX618495	-----	0
Ascomycota_sp._KX618493	-----	0
Ascomycota_sp._KX618494	-----	0

Epicoccum_nigrum_MF509753	-----	0
Boeremia_exigua_KX618486	-----	0
Boeremia_exigua_KX618484	-----	0
Boeremia_exigua_KX618485	-----	0
Cryptosphaeria_eunomia_KX618490	-----	0
Cryptosphaeria_eunomia_KX618491	-----	0
Phomopsis_sp._MF509748	-----	0
Diaporthe_eres_MF509750	-----	0
Phomopsis_oblonga_KX618489	-----	0
Bionectria_ochroleuca_MF509749	-----	0
Fusarium_oxysporum_KX618492	-----	0
Fusarium_acumnatum_MF509746	-----	0
Fusarium_avenaceum_MF509747	-----	0
Hymenoscyphus_fraxineus_KY613994	-----	0
Diplodia_mutila_KX618487	-----	0
Diplodia_mutila_KX618488	-----	0
Paraphaeosphaeria_neglecta_MF509754	gggtacggtaacagttgaagagatgacccggccagccggtgaaatgggcatccgcagcga	240
Alternaria_alternata_MF509751	-----	0
Alternaria_sp._MF509752	-----	0
Pleosporales_sp._KX618495	-----	0
Ascomycota_sp._KX618493	-----	0
Ascomycota_sp._KX618494	-----	0
Epicoccum_nigrum_MF509753	-----	0
Boeremia_exigua_KX618486	-----	0
Boeremia_exigua_KX618484	-----	0
Boeremia_exigua_KX618485	-----	0
Cryptosphaeria_eunomia_KX618490	-----	0
Cryptosphaeria_eunomia_KX618491	-----	0
Phomopsis_sp._MF509748	-----	0
Diaporthe_eres_MF509750	-----	0
Phomopsis_oblonga_KX618489	-----	0
Bionectria_ochroleuca_MF509749	-----	0
Fusarium_oxysporum_KX618492	-----	0
Fusarium_acumnatum_MF509746	-----	0
Fusarium_avenaceum_MF509747	-----	0
Hymenoscyphus_fraxineus_KY613994	-----	0
Diplodia_mutila_KX618487	-----	0
Diplodia_mutila_KX618488	-----	0
Paraphaeosphaeria_neglecta_MF509754	agccccttctgttggtgtagcagccttgccgcagccaaacatggggaacgttcacaga	300
Alternaria_alternata_MF509751	-----	0
Alternaria_sp._MF509752	-----	0
Pleosporales_sp._KX618495	-----	0
Ascomycota_sp._KX618493	-----	0
Ascomycota_sp._KX618494	-----	0
Epicoccum_nigrum_MF509753	-----	0
Boeremia_exigua_KX618486	-----	0
Boeremia_exigua_KX618484	-----	0
Boeremia_exigua_KX618485	-----	0
Cryptosphaeria_eunomia_KX618490	-----	0
Cryptosphaeria_eunomia_KX618491	-----	0
Phomopsis_sp._MF509748	-----	0
Diaporthe_eres_MF509750	-----	0
Phomopsis_oblonga_KX618489	-----	0
Bionectria_ochroleuca_MF509749	-----	0
Fusarium_oxysporum_KX618492	-----	0
Fusarium_acumnatum_MF509746	-----	0
Fusarium_avenaceum_MF509747	-----	0
Hymenoscyphus_fraxineus_KY613994	-----	0
Diplodia_mutila_KX618487	-----	0
Diplodia_mutila_KX618488	-----	0
Paraphaeosphaeria_neglecta_MF509754	ctaagtgggaagtggtggggcctagccctgcttaagatatagtgggcccccttggaac	360
Alternaria_alternata_MF509751	-----	0
Alternaria_sp._MF509752	-----tc	2
Pleosporales_sp._KX618495	-----	0
Ascomycota_sp._KX618493	-----	0
Ascomycota_sp._KX618494	-----	0
Epicoccum_nigrum_MF509753	-----	0
Boeremia_exigua_KX618486	-----	0

Boeremia_exigua_KX618484	-----	0
Boeremia_exigua_KX618485	-----	0
Cryptosphaeria_eunomia_KX618490	-----ttccgtaggtgaacct	16
Cryptosphaeria_eunomia_KX618491	-----ttccgtaggtgaacct	16
Phomopsis_sp._MF509748	-tcttgggtcatttagaggaagtaa-----aagtcgtaacaaggtctccgtaggtgaacca	54
Diaporthe_eres_MF509750	-tcttgggtcatttagaggaagtaa-----aagtcgtaacaaggtctccgtaggtgaacca	54
Phomopsis_oblonga_KX618489	-----aacct	5
Bionectria_ochroleuca_MF509749	--cttgggtcatttagaggaagtaa-----aagtcgtaacaaggtctccgtaggtgaacca	53
Fusarium_oxysporum_KX618492	-----ttccgtaggtgaacct	16
Fusarium_acumnatum_MF509746	--cttgggtcatttagaggaagtaa-----aagtcgtaacaaggtctccgtaggtgaacca	53
Fusarium_avenaceum_MF509747	--cttgggtcatttagaggaagtaa-----aagtcgtaacaaggtctccgtaggtgaacca	53
Hymenoscyphus_fraxineus_KY613994	-----ttccgtaggtgaacct	16
Diplodia_mutila_KX618487	-----ccttccgtaggtgaacct	18
Diplodia_mutila_KX618488	-----cttccgtaggtgaacct	17
Paraphaeosphaeria_neglecta_MF509754	ttgggggggtgagttcggcctcctggcggctccaagcaaaccttccgtaggtgaacct	420
Alternaria_alternata_MF509751	-----gaggaagtaa-----aagtcgtaacaaggtctccgtaggtgaacct	41
Alternaria_sp._MF509752	ttcttgggtcatttagaggaagtaa-----aagtcgtaacaaggtctccgtaggtgaacct	57
Pleosporales_sp._KX618495	-----ttccgtaggtgaacct	17
Ascomycota_sp._KX618493	-----ccgtaggtgaacct	14
Ascomycota_sp._KX618494	-----cttccgtaggtgaacct	17
Epicoccum_nigrum_MF509753	-tcttgggtcatttagaggaagtaa-----aagtcgtaacaaggttccgtaggtgaacct	54
Boeremia_exigua_KX618486	-----gaacct	6
Boeremia_exigua_KX618484	-----cttccgtaggtgaacct	17
Boeremia_exigua_KX618485	-----ccgtaggtgaacct	14
	****	
Cryptosphaeria_eunomia_KX618490	gcggagggatcattacagagttatccaactcccaaacccatgtgaacctacctatgttgc	76
Cryptosphaeria_eunomia_KX618491	gcggagggatcattacagagttatccaactcccaaacccatgtgaacctacctatgttgc	76
Phomopsis_sp._MF509748	gcggagggatcattgctggaacgcgcccagggcgaccaccagaaaaccccttctgtaacctat	114
Diaporthe_eres_MF509750	gcggagggatcattgctggaacgcgcccagggcgaccaccagaaaaccccttctgtaacctat	114
Phomopsis_oblonga_KX618489	gcggagggatcattgctggaacgcgcccagggcgaccaccagaaaaccccttctgtaacctat	65
Bionectria_ochroleuca_MF509749	gcggagggatcattaccaggtttacaactcccaaacccatgtgaacatacctactgttgc	113
Fusarium_oxysporum_KX618492	gcggagggatcattaccaggtttacaactcccaaacccctgtgaacataccttaagtgtg	76
Fusarium_acumnatum_MF509746	gcggagggatcattaccaggtttacaactcccaaacccctgtgaacataccttaagtgtg	113
Fusarium_avenaceum_MF509747	gcggagggatcattaccaggtttacaactcccaaacccctgtgaacataccttaagtgtg	113
Hymenoscyphus_fraxineus_KY613994	gcggaaggatcattacagagttcctgcccctcaagggtagaacccc-----	62
Diplodia_mutila_KX618487	gcggaaggatcattaccaggttctcgagcttcggcctcgaatctcccaccccttctgtaaca	78
Diplodia_mutila_KX618488	gcggaaggatcattaccaggttctcgagcttcggcctcgaatctcccaccccttctgtaaca	77
Paraphaeosphaeria_neglecta_MF509754	gcggaaggatcattatccatctcaaacaggtgcggctcgccggcccccggggcttgcctc	480
Alternaria_alternata_MF509751	gcggagggatcattaccaggtttacaactcccaaacccctgtgaacctct-----	84
Alternaria_sp._MF509752	gcggagggatcattaccaggtttacaactcccaaacccctgtgaacctct-----	100
Pleosporales_sp._KX618495	gcggaaggatcattatccttttctcgagggttgatgccaatgctctgggctcatgctcg	77
Ascomycota_sp._KX618493	gcggaaggatcattaccaa-----tt	35
Ascomycota_sp._KX618494	gcggaaggatcattaccaa-----tt	38
Epicoccum_nigrum_MF509753	gcggaaggatcattaccta-----ga	75
Boeremia_exigua_KX618486	gcggaaggatcattaccta-----ga	27
Boeremia_exigua_KX618484	gcggaaggatcattaccta-----ga	38
Boeremia_exigua_KX618485	gcggaaggatcattaccta-----ga	35
	*****	
Cryptosphaeria_eunomia_KX618490	ctcggcggggaagcct-----accggtagctaccctgtagctaccgggagcgagct	129
Cryptosphaeria_eunomia_KX618491	ctcggcggggaagcct-----accggtagctaccctgtagctaccgggagcgagct	129
Phomopsis_sp._MF509748	acctactgttgccctc-----ggcg-----catgccggcccccttggggcc	156
Diaporthe_eres_MF509750	acctactgttgccctc-----ggcg-----catgccggcccccttggggcc	156
Phomopsis_oblonga_KX618489	acctactgttgccctc-----ggcg-----ctagctg--gtccctcggggcc	105
Bionectria_ochroleuca_MF509749	ttcggcgggattgcc-----ccg-----gg	133
Fusarium_oxysporum_KX618492	cctcggcggatcagc-----ccg-----cg	96
Fusarium_acumnatum_MF509746	cctcggcggatcagc-----ccg-----cg	133
Fusarium_avenaceum_MF509747	cctcggcggatcagc-----ccg-----cg	133
Hymenoscyphus_fraxineus_KY613994	-----accctgtgtatattatattgttgccttagcaggtcg	99
Diplodia_mutila_KX618487	ta-----cctctgttgccttggcggctcttgcggctggaggcc	117
Diplodia_mutila_KX618488	ta-----cctctgttgccttggcggctcttgcggctggaggcc	116
Paraphaeosphaeria_neglecta_MF509754	cggggtgtaggggtaaacctgcgcggcggcctgtaccctcttttacagcaacc	540
Alternaria_alternata_MF509751	-cggggttaca-----gcctgtgtaattattacccttcttcttgcgta--	129
Alternaria_sp._MF509752	-cggggttaca-----gcctgtgtaattattacccttcttcttgcgta--	145
Pleosporales_sp._KX618495	gttactggct-----catctcttctgattctaccatgtttttgcgta--	123
Ascomycota_sp._KX618493	taatcggggga-----cttccgctcctgtctgaaccttcttcttgcgta--	81
Ascomycota_sp._KX618494	taatcggggga-----cttccgctcctgtctgaaccttcttcttgcgta--	84
Epicoccum_nigrum_MF509753	gtttagtagct-----ctcggctcgtacctcttaccatgtcttttagta--	121
Boeremia_exigua_KX618486	gtttagtagct-----tg--cctaccatctcttaccatgtcttttagta--	71
Boeremia_exigua_KX618484	gtttagtagct-----tg--cctaccatctcttaccatgtcttttagta--	82
Boeremia_exigua_KX618485	gtttagtagct-----tg--cctaccatctcttaccatgtcttttagta--	79

Cryptosphaeria_eunomia_KX618490	accctgtagcc---cgctgcaggcctaccgccgggtggacgattaaactcttgtttttta	186
Cryptosphaeria_eunomia_KX618491	accctgtagcc---cgctgcaggcctaccgccgggtggacgattaaactcttgtttttta	186
Phomopsis_sp._MF509748	cctctcctg-----gaggagcagcagccggcggccaactcttgtttttac	208
Diaporthe_eres_MF509750	cctcacccccc---ggtgttgagacagcccgccggcggccaactcttgtttttac	213
Phomopsis_oblonga_KX618489	cctcacctctg---ggtgttgagacagcccgccggcggccaactcttgtttttac	162
Bionectria_ochroleuca_MF509749	cgctctgtgtg---ccccgatcaggcgccgcctaggaacttaattcttgtttttatt	190
Fusarium_oxysporum_KX618492	ccccgtaaa-----acgggacggcccgccagaggaccaccaactctaattgtttctta	147
Fusarium_acumnatum_MF509746	ccccgtaaa-----acgggacggcccgccagaggaccaccaactctaattgtttctta	184
Fusarium_avenaceum_MF509747	ccccgtaaa-----acgggacggcccgccagaggaccaccaactctaattgtttctta	184
Hymenoscyphus_fraxineus_KY613994	c---cctctgggcgtcggcctcggc-tgactgtgctgctagaggaccctaaattttga	154
Diplodia_mutila_KX618487	ctcaaaaagccccc---gtcgcc-ttccgcccagaggaccttcaactccagtcagta	172
Diplodia_mutila_KX618488	ctcaaaaagccccc---gtcgcc-ttccgcccagaggaccttcaactccagtcagta	171
Paraphaeosphaeria_neglecta_MF509754	tttcgtctcctctcgccggggcaactcgccgctggaacaaaaaaccctttttgtcat	600
Alternaria_alternata_MF509751	cttctgtttcctctggtgggttcgcccacca---ctaggacaacataaacctttgtga	185
Alternaria_sp._MF509752	ctctctgtttcctctggtgggttcgcccacca---ctaggacaacataaacctttgtga	201
Pleosporales_sp._KX618495	ctacttgtttcctctggtggccttaccac---ataggacatattaaaaccctttgtga	179
Ascomycota_sp._KX618493	ctagtgtttcctcctggtaggcttgcctgccg---gttggacagtaaaaaacccttt-gt	137
Ascomycota_sp._KX618494	ctagtgtttcctcctggtaggcttgcctgccg---gttggacagtaaaaaacccttt-gt	140
Epicoccum_nigrum_MF509753	cct-tcgtttcctcctggtgggtccgcccgccg---attggacaacattcaaacctttgca	177
Boeremia_exigua_KX618486	cct-tcgtttcctcctggtgggtccgcccgccg---attgga-caaacttaaacctttgta	126
Boeremia_exigua_KX618484	cct-tcgtttcctcctggtgggtccgcccgccg---attgga-caaacttaaacctttgta	137
Boeremia_exigua_KX618485	cct-tcgtttcctcctggtgggtccgcccgccg---attgga-caaacttaaacctttgta	134
	*	
Cryptosphaeria_eunomia_KX618490	gtggattatctgagtgtttatacttaataagttaaaactttcaacaacggatctcttgg	246
Cryptosphaeria_eunomia_KX618491	gtggattatctgagtgtttatacttaataagttaaaactttcaacaacggatctcttgg	246
Phomopsis_sp._MF509748	actgaaactctgagataaaaataaataagatcaaaaactttcaacaacggatctcttgg	268
Diaporthe_eres_MF509750	actgaaactctgagataaaaataaataagatcaaaaactttcaacaacggatctcttgg	273
Phomopsis_oblonga_KX618489	actgaaactctgagataaaaataaataagatcaaaaactttcaacaacggatctcttgg	222
Bionectria_ochroleuca_MF509749	tggaaactctctgagtgttttcaaaaataaataaactttcaacaacggatctcttgg	250
Fusarium_oxysporum_KX618492	ttgtaactctgagtaaaaacaaataaataaactttcaacaacggatctcttgg	207
Fusarium_acumnatum_MF509746	ttgtaactctgagtaaaaacaaataaataaactttcaacaacggatctcttgg	244
Fusarium_avenaceum_MF509747	ttgtaactctgagtaaaaacaaataaataaactttcaacaacggatctcttgg	244
Hymenoscyphus_fraxineus_KY613994	aatacagtcgtcgtctgagtactatttaagttaaaactttcaacaacggatctcttgg	214
Diplodia_mutila_KX618487	aacgtcgcagtcgtgaaaacaagttataaactaaaactttcaacaacggatctcttgg	232
Diplodia_mutila_KX618488	aacgtcgcagtcgtgaaaacaagttataaactaaaactttcaacaacggatctcttgg	231
Paraphaeosphaeria_neglecta_MF509754	ctagcatctctgctgatacaaaaactctgttacaactttcaacaatggatctcttgg	660
Alternaria_alternata_MF509751	attgcaatcagcgtcagtaacaaataaataaactcaactttcaacaacggatctcttgg	245
Alternaria_sp._MF509752	attgcaatcagcgtcagtaacaaataaataaactcaactttcaacaacggatctcttgg	261
Pleosporales_sp._KX618495	attgcagtcagcgtcagtaacaaataaataaactcaactttcaacaacggatctcttgg	239
Ascomycota_sp._KX618493	aattgcaatcagcgtcagaaaaacataaataagttacaactttcaacaacggatctcttgg	197
Ascomycota_sp._KX618494	aattgcaatcagcgtcagaaaaacataaataagttacaactttcaacaacggatctcttgg	200
Epicoccum_nigrum_MF509753	gttgcaatcagcgtcgtgaaaaacataaataagttacaactttcaacaacggatctcttgg	237
Boeremia_exigua_KX618486	attgaaatcagcgtcgtgaaaaacataaataagttacaactttcaacaacggatctcttgg	186
Boeremia_exigua_KX618484	attgaaatcagcgtcgtgaaaaacataaataagttacaactttcaacaacggatctcttgg	197
Boeremia_exigua_KX618485	attgaaatcagcgtcgtgaaaaacataaataagttacaactttcaacaacggatctcttgg	194
	* * * * *	
Cryptosphaeria_eunomia_KX618490	tctggcatcgatgaagaacgcagcgaatgcgataagtaattggaattgcagaattcagt	306
Cryptosphaeria_eunomia_KX618491	tctggcatcgatgaagaacgcagcgaatgcgataagtaattggaattgcagaattcagt	306
Phomopsis_sp._MF509748	tctggcatcgatgaagaacgcagcgaatgcgataagtaattggaattgcagaattcagt	328
Diaporthe_eres_MF509750	tctggcatcgatgaagaacgcagcgaatgcgataagtaattggaattgcagaattcagt	333
Phomopsis_oblonga_KX618489	tctggcatcgatgaagaacgcagcgaatgcgataagtaattggaattgcagaattcagt	282
Bionectria_ochroleuca_MF509749	tctggcatcgatgaagaacgcagcgaatgcgataagtaattggaattgcagaattcagt	310
Fusarium_oxysporum_KX618492	tctggcatcgatgaagaacgcagcgaatgcgataagtaattggaattgcagaattcagt	267
Fusarium_acumnatum_MF509746	tctggcatcgatgaagaacgcagcgaatgcgataagtaattggaattgcagaattcagt	304
Fusarium_avenaceum_MF509747	tctggcatcgatgaagaacgcagcgaatgcgataagtaattggaattgcagaattcagt	304
Hymenoscyphus_fraxineus_KY613994	tctggcatcgatgaagaacgcagcgaatgcgataagtaattggaattgcagaattcagt	274
Diplodia_mutila_KX618487	tctggcatcgatgaagaacgcagcgaatgcgataagtaattggaattgcagaattcagt	292
Diplodia_mutila_KX618488	tctggcatcgatgaagaacgcagcgaatgcgataagtaattggaattgcagaattcagt	291
Paraphaeosphaeria_neglecta_MF509754	tctggcatcgatgaagaacgcagcgaatgcgataagtaattggaattgcagaattcagt	720
Alternaria_alternata_MF509751	tctggcatcgatgaagaacgcagcgaatgcgataagtaattggaattgcagaattcagt	305
Alternaria_sp._MF509752	tctggcatcgatgaagaacgcagcgaatgcgataagtaattggaattgcagaattcagt	321
Pleosporales_sp._KX618495	tctggcatcgatgaagaacgcagcgaatgcgataagtaattggaattgcagaattcagt	299
Ascomycota_sp._KX618493	tctggcatcgatgaagaacgcagcgaatgcgataagtaattggaattgcagaattcagt	257
Ascomycota_sp._KX618494	tctggcatcgatgaagaacgcagcgaatgcgataagtaattggaattgcagaattcagt	260
Epicoccum_nigrum_MF509753	tctggcatcgatgaagaacgcagcgaatgcgataagtaattggaattgcagaattcagt	297
Boeremia_exigua_KX618486	tctggcatcgatgaagaacgcagcgaatgcgataagtaattggaattgcagaattcagt	246
Boeremia_exigua_KX618484	tctggcatcgatgaagaacgcagcgaatgcgataagtaattggaattgcagaattcagt	257
Boeremia_exigua_KX618485	tctggcatcgatgaagaacgcagcgaatgcgataagtaattggaattgcagaattcagt	254
	*****	
Cryptosphaeria_eunomia_KX618490	gaatcactcgaatctttgaacgcacattgcccattagatctctaggcctgctgtt	366

Cryptosphaeria_eunomia_KX618491	gaatcatcgaatccttgaacgcacattgcccatttagtattctagtgggcatgcctggt	366
Phomopsis_sp._MF509748	gaatcatcgaatccttgaacgcacattgcccctctggtattccggagggcatgcctggt	388
Diaporthe_eres_MF509750	gaatcatcgaatccttgaacgcacattgcccctctggtattccggagggcatgcctggt	393
Phomopsis_oblonga_KX618489	gaatcatcgaatccttgaacgcacattgcccctctggtattccggagggcatgcctggt	342
Bionectria_ochroleuca_MF509749	gaatcatcgaatccttgaacgcacattgcccgcagattctgctgggcatgcctggt	370
Fusarium_oxysporum_KX618492	gaatcatcgaatccttgaacgcacattgcccgcctggtattccgggggcatgcctggt	327
Fusarium_acumnatum_MF509746	gaatcatcgaatccttgaacgcacattgcccgcctggtattccgggggcatgcctggt	364
Fusarium_avenaceum_MF509747	gaatcatcgaatccttgaacgcacattgcccgcctggtattccgggggcatgcctggt	364
Hymenoscyphus_fraxineus_KY613994	gaatcatcgaatccttgaacgcacattgcccgcctggtattccgggggcatgcctggt	334
Diplodia_mutila_KX618487	gaatcatcgaatccttgaacgcacattgcccgccttggcattccgaggggcatgcctggt	352
Diplodia_mutila_KX618488	gaatcatcgaatccttgaacgcacattgcccgccttggcattccgaggggcatgcctggt	351
Paraphaeosphaeria_neglecta_MF509754	gaatcatcgaatccttgaacgcacattgcccgccttgggtattccatgggggcatgcctggt	780
Alternaria_alternata_MF509751	gaatcatcgaatccttgaacgcacattgcccgccttgggtattccaaagggcatgcctggt	365
Alternaria_sp._MF509752	gaatcatcgaatccttgaacgcacattgcccgccttgggtattccaaagggcatgcctggt	381
Pleosporales_sp._KX618495	gaatcatcgaatccttgaacgcacattgcccgccttgggtattccatggggcatgcctggt	359
Ascomycota_sp._KX618493	gaatcatcgaatccttgaacgcacattgcccgccttgggtattccatggggcatgcctggt	317
Ascomycota_sp._KX618494	gaatcatcgaatccttgaacgcacattgcccgccttgggtattccatgggggcatgcctggt	320
Epicoccum_nigrum_MF509753	gaatcatcgaatccttgaacgcacattgcccgccttgggtattccatgggggcatgcctggt	357
Boeremia_exigua_KX618486	gaatcatcgaatccttgaacgcacattgcccgccttgggtattccatgggggcatgcctggt	306
Boeremia_exigua_KX618484	gaatcatcgaatccttgaacgcacattgcccgccttgggtattccatgggggcatgcctggt	317
Boeremia_exigua_KX618485	gaatcatcgaatccttgaacgcacattgcccgccttgggtattccatgggggcatgcctggt	314
	***** * *** *****	
Cryptosphaeria_eunomia_KX618490	cgagcgtcatttcgaccattcaagccctgggtgcttgggtggtgggagcttgcctc----	422
Cryptosphaeria_eunomia_KX618491	cgagcgtcatttcgaccattcaagccctgggtgcttgggtggtgggagcttgcctc----	422
Phomopsis_sp._MF509748	cgagcgtcatttcaaccctcaagcctggcttgggtgatggggcactgcctgtaa-----	441
Diaporthe_eres_MF509750	cgagcgtcatttcaaccctcaagcctggcttgggtgatggggcactgccttacc-----	449
Phomopsis_oblonga_KX618489	cgagcgtcatttcaaccctcaagcctggcttgggtgatggggcactgccttacc-----	398
Bionectria_ochroleuca_MF509749	tgagcgtcatttcaaccctcatgcccctaggcctgggttgggggatcgccaaagcccg	430
Fusarium_oxysporum_KX618492	cgagcgtcatttcaaccctcaagcccccgggttgggttggggat-cggctctgc--ct	384
Fusarium_acumnatum_MF509746	cgagcgtcatttcaaccctcaagcccccgggttgggttggggat-cggctctgc--ct	423
Fusarium_avenaceum_MF509747	cgagcgtcatttcaaccctcaagcccccgggttgggttggggat-cggctctgc--cct	421
Hymenoscyphus_fraxineus_KY613994	cgagcgtcatttagaccactcccctcggcggggtcttgggctgcgccttggggggg	394
Diplodia_mutila_KX618487	cgagcgtcatttcaaccctcaagctctgcttgggtattggggcagctcctctcgcgagc	412
Diplodia_mutila_KX618488	cgagcgtcatttcaaccctcaagctctgcttgggtattggggcagctcctctcgcgagc	411
Paraphaeosphaeria_neglecta_MF509754	cgagcgtcatttcaaccctcaagctctgcttgggtattggggcagctcctctcgcgagc	840
Alternaria_alternata_MF509751	cgagcgtcattttagaccctcaagcttgccttgggttggggcgtctgtctctagct----	421
Alternaria_sp._MF509752	cgagcgtcattttagaccctcaagcttgccttgggttggggcgtctgtctctagct----	437
Pleosporales_sp._KX618495	cgagcgtcattttagaccctcaagcttgccttgggttggggcgtctgtctctagct----	419
Ascomycota_sp._KX618493	cgagcgtcattttagaccctcaagctctgcttgggttgggggttgggtcttgcctcctg	377
Ascomycota_sp._KX618494	cgagcgtcattttagaccctcaagctctgcttgggttgggggttgggtcttgcctcctg	380
Epicoccum_nigrum_MF509753	cgagcgtcattttagaccctcaagctctgcttgggttgggggttgggtctcgcctctg	417
Boeremia_exigua_KX618486	cgagcgtcattttagaccctcaagctctgcttgggttgggggttgggtctcgcctctg	366
Boeremia_exigua_KX618484	cgagcgtcattttagaccctcaagctctgcttgggttgggggttgggtctcgcctctg	377
Boeremia_exigua_KX618485	cgagcgtcattttagaccctcaagctctgcttgggttgggggttgggtctcgcctctg	374
	***** **	
Cryptosphaeria_eunomia_KX618490	-----ggacaactcctcaaaagcattggcggagtcgcggtggcccca--agcgtag	471
Cryptosphaeria_eunomia_KX618491	-----ggacaactcctcaaaagcattggcggagtcgcggtggcccca--agcgtag	471
Phomopsis_sp._MF509748	-----aa--ggcagggccctgaaatcagtgggcagctcggcaggaaccccg--agcgcag	492
Diaporthe_eres_MF509750	----aag--gagcagggcctgaaatcagtgggcagctcggcaggaaccccg--agcgcag	501
Phomopsis_oblonga_KX618489	----aag--aagcagggccctgaaatcagtgggcagctcggcaggaaccccg--agcgcag	450
Bionectria_ochroleuca_MF509749	cgagggga--cggccggccctaaatctagtggcggaccgctcgtggcctcctctgcgga	488
Fusarium_oxysporum_KX618492	cagcggc--gtgcccggcccgaaatacatggcggtctcgtgcagcctccattgctgtag	442
Fusarium_acumnatum_MF509746	ctgggcg--gtgcccggcccgaaatacatggcggtctcgtgcagcctccattgctgtag	481
Fusarium_avenaceum_MF509747	tctggcg--gtgcccggcccgaaatacatggcggtctcgtgcagcctccattgctgtag	479
Hymenoscyphus_fraxineus_KY613994	ccttaaac-----cagtgggcggtgcctaaagctcctacgcgtagttaa	437
Diplodia_mutila_KX618487	cgccctcaaaagcctcggcggtggctgttcagccctcaagcgtagttag-----aatacac	466
Diplodia_mutila_KX618488	cgccctcaaaagcctcggcggtggctgttcagccctcaagcgtagttag-----aatacac	465
Paraphaeosphaeria_neglecta_MF509754	-----gcgactgcgcccaaatggcagcggcttgcctcctc--cgcgca	891
Alternaria_alternata_MF509751	---ttgctggagactcgccttaaagtaattggcagccggcctactggttccggagcgcag	478
Alternaria_sp._MF509752	---ttgctggagactcgccttaaagtaattggcagccggcctactggttccggagcgcag	494
Pleosporales_sp._KX618495	gcgtaagtggagactcgccttaaaacaattggcagccggcctactggttccggagcgcag	478
Ascomycota_sp._KX618493	-----gcgactcgcctcaaaacaattggcagccggcctactggttccggagcgcag	429
Ascomycota_sp._KX618494	-----gcgactcgcctcaaaacaattggcagccggcctactggttccggagcgcag	432
Epicoccum_nigrum_MF509753	-----gtagactcgccttaaaacaattggcagccggcctactggttccggagcgcag	469
Boeremia_exigua_KX618486	-----gtagactcgccttaaaacaattggcagccggcctactggttccggagcgcag	418
Boeremia_exigua_KX618484	-----gtagactcgccttaaaacaattggcagccggcctactggttccggagcgcag	429
Boeremia_exigua_KX618485	-----gtagactcgccttaaaacaattggcagccggcctactggttccggagcgcag	426
	* * *	
Cryptosphaeria_eunomia_KX618490	taattcttctcgttccggtgtgtcagcggtggcgtctggccgttaaaccctccatttt	531
Cryptosphaeria_eunomia_KX618491	taattcttctcgttccggtgtgtcagcggtggcgtctggccgttaaaccctccatttt	531

Phomopsis_sp._MF509748	tagttaaacctcgctc--tggaaggccctggcggtgccctgccgttaaaccocccaactt	550
Diaporthe_eres_MF509750	tagttaaacctcgctc--tggaaggccctggcggtgccctgccgttaaaccocccaactt	559
Phomopsis_oblonga_KX618489	tagttaaacctcgctc--tggaaggccctggcggtgccctgccgttaaaccocccaactt	508
Bionectria_ochroleuca_MF509749	tagtgatattccgcat-----cggagagcgagccctgccctgccgttaaaccocccaactt	542
Fusarium_oxysporum_KX618492	tagctaaccctcgca-----actggaacgcgcgcgccatgccgtaaaaccocccaactt	496
Fusarium_acumnatum_MF509746	tagctaaccctcgca-----actggaacgcgcgcgccatgccgtaaaaccocccaactt	535
Fusarium_avenaceum_MF509747	tagctaaccctcgca-----actggaacgcgcgcgccatgccgtaaaaccocccaactt	533
Hymenoscyphus_fraxineus_KY613994	tt-----cttctcgcg--atagggctcctggcggtgtcttggcagcaaccocccaactt	486
Diplodia_mutila_KX618487	ct-----cgcttggagcggttg-gcgtcgcccgccggagcaacccttctgaacttt	516
Diplodia_mutila_KX618488	ct-----cgcttggagcggttg-gcgtcgcccgccggagcaacccttctgaacttt	515
Paraphaeosphaeria_neglecta_MF509754	ca-----caattgcgctctcgcg--ggggcgctggcccgctccacggaacaactt	939
Alternaria_alternata_MF509751	ca-----caagtgcactctctatcagcaaaaggtctagcatccattaagccttttt	529
Alternaria_sp._MF509752	ca-----caagtgcactctctatcagcaaaaggtctagcatccattaagccttttt	545
Pleosporales_sp._KX618495	ca-----cattttgcgctcttgtca-tga-ttgttgcatccatcaagccttttt	527
Ascomycota_sp._KX618493	ca-----cattttgcgccccctcagctttaa-cgcttgcgctccattaaagctcaactt	479
Ascomycota_sp._KX618494	ca-----cattttgcgccccctcagctttaa-cgcttgcgctccattaaagctcaactt	482
Epicoccum_nigrum_MF509753	ta-----catctcgcttcttgactcaca-cgacga--cgtccaaaagctacattt	517
Boeremia_exigua_KX618486	ta-----catctcgcttcttgactcaca-cgacga--cgtccaaaagctacattt	466
Boeremia_exigua_KX618484	ta-----catctcgcttcttgactcaca-cgacga--cgtccaaaagctacattt	477
Boeremia_exigua_KX618485	ta-----catctcgcttcttgactcaca-cgacga--cgtccaaaagctacattt	474
	*	
Cryptosphaeria_eunomia_KX618490	tcagtgggttgacctcggatcaggttaggaataaccgctgaacttaagcatatcaataaagc	591
Cryptosphaeria_eunomia_KX618491	tcagtgggttgacctcggatcaggttaggaataaccgctgaacttaagcatatcaataaagc	588
Phomopsis_sp._MF509748	ctgaaaatttgacctcggatcaggttaggaataaccgctgaacttaagcatatcaataaagc	610
Diaporthe_eres_MF509750	ctgaaaatttgacctcggatcaggttaggaataaccgctgaacttaagcatatcaataaagc	619
Phomopsis_oblonga_KX618489	ctgaaaatttgacctcggatcaggttaggaataaccgctgaacttaagcatatcaataaagc	565
Bionectria_ochroleuca_MF509749	ttccaagggttgacctcagctcaggttaggaataaccgctgaacttaagcatatcaataaagc	602
Fusarium_oxysporum_KX618492	tctgaatggttgacctcggatcaggttaggaataaccgctgaacttaagcatatcaataaagc	556
Fusarium_acumnatum_MF509746	tctgaatggttgacctcggatcaggttaggaataaccgctgaacttaagcatatcaataaagc	595
Fusarium_avenaceum_MF509747	tctgaatggttgacctcggatcaggttaggaataaccgctgaacttaagcatatcaataaagc	593
Hymenoscyphus_fraxineus_KY613994	ctctagggttgacctcggatcaggttaggaataaccgctgaacttaagcatatcaataaagc	546
Diplodia_mutila_KX618487	tctcaagggttgacctcggatcaggttaggaataaccgctgaacttaagcatatcaataaagc	573
Diplodia_mutila_KX618488	tctcaagggttgacctcggatcaggttaggaataaccgctgaacttaagcatatcaataaagc	575
Paraphaeosphaeria_neglecta_MF509754	taccgtctttgacctcggatcaggttaggaataaccgctgaacttaagcatatcaataaagc	994
Alternaria_alternata_MF509751	ttcaacttttgacctcggatcaggttaggaataaccgctgaacttaagcatatcaataaagc	585
Alternaria_sp._MF509752	ttcaacttttgacctcggatcaggttaggaataaccgctgaacttaagcatatcaataaagc	605
Pleosporales_sp._KX618495	attagctcttgacctcggatcaggttaggaataaccgctgaacttaagcatatcaataaagc	587
Ascomycota_sp._KX618493	ttttgctcttgacctcggatcaggttaggaataaccgctgaacttaagcatatcaataaagc	539
Ascomycota_sp._KX618494	ttttgctcttgacctcggatcaggttaggaataaccgctgaacttaagcatatcaataaagc	542
Epicoccum_nigrum_MF509753	ttacactcttgacctcggatcaggttaggaataaccgctgaacttaagcatatcaataaagc	577
Boeremia_exigua_KX618486	ttacactcttgacctcggatcaggttaggaataaccgctgaacttaagcatatcaataaagc	526
Boeremia_exigua_KX618484	ttacactcttgacctcggatcaggttaggaataaccgctgaacttaagcatatcaataaagc	537
Boeremia_exigua_KX618485	ttacactcttgacctcggatcaggttaggaataaccgctgaacttaagcatatcaataaagc	531
	*****	
Cryptosphaeria_eunomia_KX618490	ggagga-	597
Cryptosphaeria_eunomia_KX618491	-----	588
Phomopsis_sp._MF509748	ggaggaa	617
Diaporthe_eres_MF509750	ggaggaa	626
Phomopsis_oblonga_KX618489	-----	565
Bionectria_ochroleuca_MF509749	ggagga-	608
Fusarium_oxysporum_KX618492	ggagga-	562
Fusarium_acumnatum_MF509746	ggaggaa	602
Fusarium_avenaceum_MF509747	ggaggaa	600
Hymenoscyphus_fraxineus_KY613994	ggagga-	552
Diplodia_mutila_KX618487	-----	573
Diplodia_mutila_KX618488	ggagga-	581
Paraphaeosphaeria_neglecta_MF509754	-----	994
Alternaria_alternata_MF509751	-----	585
Alternaria_sp._MF509752	ggaggaa	612
Pleosporales_sp._KX618495	ggaggaa	594
Ascomycota_sp._KX618493	ggagga-	545
Ascomycota_sp._KX618494	ggagga-	548
Epicoccum_nigrum_MF509753	ggaggaa	584
Boeremia_exigua_KX618486	ggagga-	532
Boeremia_exigua_KX618484	ggaggaa	544
Boeremia_exigua_KX618485	-----	531